

Kinetic Monte Carlo Simulations of Surface Reactions

Michail Stamatakis

University College London

Kinetic Monte Carlo Crash-Course and Workshop

<http://zacros.org/workshop>


Why Do Kinetic Modelling in Catalysis and Surface Science? ²

- **DFT Calculations in Catalysis:**



- ✓ Electronic structure of materials
- ✓ Stability of intermediates
- ✓ Chemical pathways and energy barriers

*Need “predictive models able to capture trends in **activity** and **selectivity**”*




- Parallel or competing pathways?
- Temperature and pressure effects?
- Coverage effects on reaction rates?



Kinetic modelling necessary



The Kinetic Monte Carlo Approach






- Instead of simulating dynamics, KMC¹ focuses on rare events
- Simulates reactions much faster than Molecular Dynamics
- Incorporates spatial information contrary to micro-kinetic models²

¹ M. Neurock and E. W. Hansen, *Comput. Chem. Eng.* 22, S1045 (1998); K. Reuter and M. Scheffler, *Phys. Rev. Lett.* 90: 046103 (2003); M. Stamatakis, *J. Phys. Condens. Matter.* 27: 013001 (2015).

² J. A. Dumesic et al., *The Microkinetics of Heterogeneous Catalysis.* (American Chemical Society, 1993).

The Kinetic Monte Carlo Approach




- Instead of simulating dynamics, KMC¹ focuses on rare events
- Simulates reactions much faster than Molecular Dynamics
- Incorporates spatial information contrary to micro-kinetic models²

¹ M. Neurock and E. W. Hansen, *Comput. Chem. Eng.* 22, S1045 (1998); K. Reuter and M. Scheffler, *Phys. Rev. Lett.* 90: 046103 (2003); M. Stamatakis, *J. Phys. Condens. Matter*, 27: 013001 (2015).

² J. A. Dumesic et al., *The Microkinetics of Heterogeneous Catalysis*. (American Chemical Society, 1993).


Outline

- Atomistic/Molecular level
 - Calculating rates for elementary events (transition state theory)
- Mesoscopic level
 - Simulating reactions on spatially extended systems
- Accurate modelling of catalytic surface reactions
 - Complex materials (lattices)
 - Complicated reactions
 - Coverage effects



Outline


- **Atomistic/Molecular level**
 - Calculating rates for elementary events (transition state theory)
- Mesoscopic level
 - Simulating reactions on spatially extended systems
- Accurate modelling of catalytic surface reactions
 - Complex materials (lattices)
 - Complicated reactions
 - Coverage effects



Transition State Theory

- Assumption:** quasi-equilibrium between initial state and transition state:

$$\frac{[*]}{[A] \cdot [B]} = \frac{Q_*}{Q_A \cdot Q_B}$$



$$Q_A = \frac{1}{h^{3 \cdot N_A}} \cdot \int \exp\left(-\frac{\mathcal{H}_A}{k_B \cdot T}\right) d\Gamma_A \quad \text{and} \quad Q_B = \frac{1}{h^{3 \cdot N_B}} \cdot \int \exp\left(-\frac{\mathcal{H}_B}{k_B \cdot T}\right) d\Gamma_B$$

$$Q_* = \frac{1}{h^{3 \cdot N}} \cdot \int \exp\left(-\frac{\mathcal{H}_*}{k_B \cdot T}\right) d\Gamma_* \quad \text{where } N = N_A + N_B$$


Transition State Theory

- Assumption:** at the transition state the Hamiltonian can be expressed as:

$$\mathcal{H}_* = \frac{p^2}{2 \cdot m^*} + \mathcal{H}^\ddagger$$

kinetic energy (along reaction coordinate)

“everything else”



Therefore:

$$Q_* = \frac{\delta}{h} \cdot \frac{1}{h^{3 \cdot N - 1}} \cdot \int \exp\left(-\frac{\mathcal{H}^\ddagger}{k_B \cdot T}\right) d\Gamma^\ddagger \cdot \int_{-\infty}^{\infty} \exp\left(-\frac{p^2}{2 \cdot m^* \cdot k_B \cdot T}\right) dp$$

$$Q_* = \frac{\delta}{h} \cdot Q^\ddagger \cdot \sqrt{2 \cdot \pi \cdot m^* \cdot k_B \cdot T}$$


Transition State Theory

- Now number of transitions from reactants to products per unit time is (on average):

$$r_{\text{TST}} \cdot dt = k_{\text{TST}} \cdot [A] \cdot [B] \cdot dt =$$

$$[*] \cdot \frac{\int_0^{\infty} \frac{p \cdot dt}{\delta \cdot m^*} \cdot \exp\left(-\frac{p^2}{2 \cdot m^* \cdot k_B \cdot T}\right) dp}{\int_0^{\infty} \exp\left(-\frac{p^2}{2 \cdot m^* \cdot k_B \cdot T}\right) dp} =$$

$$\frac{[*] \cdot dt \cdot \cancel{m^*} \cdot k_B \cdot T}{\delta \cdot \cancel{m^*} \cdot \sqrt{2 \cdot \pi \cdot m^* \cdot k_B \cdot T}}$$



Transition State Theory

- Collecting the relations in red boxes: $\frac{[*]}{[A] \cdot [B]} = \frac{Q_*}{Q_A \cdot Q_B}$

$$k_{\text{TST}} = \frac{[*]}{[A] \cdot [B]} \cdot \frac{k_B \cdot T}{\delta \cdot \sqrt{2 \cdot \pi \cdot m^* \cdot k_B \cdot T}} \quad Q_* = \frac{\delta}{h} \cdot Q^\ddagger \cdot \sqrt{2 \cdot \pi \cdot m^* \cdot k_B \cdot T}$$

... we end up with:

$$k_{\text{TST}} = \frac{Q^\ddagger}{Q_A \cdot Q_B} \cdot \frac{k_B \cdot T}{h}$$



Usually, one encounters a slightly modified version:

- Fudge factor κ accounting for re-crossings
- Potential energy contribution taken out of the partition functions


$$k_{\text{TST}} = \kappa \cdot \frac{k_B \cdot T}{h} \cdot \frac{q^\ddagger}{q_{\text{reac}}} \cdot \exp\left(-\frac{E^\ddagger}{k_B \cdot T}\right)$$

So What Does It All Mean?

- $r_{\text{TST}} = k_{\text{TST}} \cdot [A] \cdot [B] \cdot dt$ = average number of transitions from reactants to products per unit time. But what about the statistics?





⋮



Ensemble


- **The system is memoryless** \Rightarrow

- Waiting time for reaction events follows the exponential distribution with rate parameter r_{TST}
- Number of events in given time interval follows the Poisson distribution with rate parameter r_{TST}



Outline

- Atomistic/Molecular level
 - Calculating rates for elementary events (transition state theory)
- **Mesoscopic level**
 - **Simulating reactions on spatially extended systems**
- Accurate modelling of catalytic surface reactions
 - Complex materials (lattices)
 - Complicated reactions
 - Coverage effects



From One Reaction to Many...

- Many species on catalytic surface, many possible reaction types,

each with its own rate constant:

$$\tau_1 \sim \text{Exp}(r_{\text{TST},1})$$

$$\tau_2 \sim \text{Exp}(r_{\text{TST},2})$$

⋮


$$\tau_n \sim \text{Exp}(r_{\text{TST},n})$$

Time of occurrence of next event:

$$\tau = \min_i (\tau_i) \sim \text{Exp}\left(\sum_{i=1}^n r_{\text{TST},i}\right)$$


Event to occur:

the one with the smallest time.



From One Reaction to Many...

- Many species on catalytic surface, many possible reaction types, each with its own rate constant.



Time of occurrence of next event:

$$\tau = \min_i (\tau_i) \sim \text{Exp} \left(\sum_{i=1}^n r_{\text{TST},i} \right)$$

Event to occur:

the one with the smallest time.

⇒ we can simulate a sequence of lattice configurations and take samples

Equation governing the statistics of these configurations?


The Master Equation

Probability of configuration σ

$$\frac{dP(\sigma)}{dt} = \sum_{\sigma'} \left[W_{\sigma' \rightarrow \sigma} \cdot P(\sigma') - W_{\sigma \rightarrow \sigma'} \cdot P(\sigma) \right]$$

Probability efflux to other configurations σ'

Probability influx from other configurations σ'



The Master Equation

Probability of configuration σ


$$\frac{dP(\sigma)}{dt} = \sum_{\sigma'} \left[W_{\sigma' \rightarrow \sigma} \cdot P(\sigma') - W_{\sigma \rightarrow \sigma'} \cdot P(\sigma) \right]$$

Probability influx from other configurations σ'





Probability efflux to other configurations σ'

- Equation linear with respect to $P(\sigma)$ but **state space** often too large, e.g. for lattice gas $2^{N_{\text{sites}}} = 5.62 \times 10^{14}$ for $N_{\text{sites}} = 49$
- ✓ We simulate & sample stochastic trajectories (KMC method)

KMC Algorithm Flowchart




Typical KMC Output






Outline

- Atomistic/Molecular level
 - Calculating rates for elementary events (transition state theory)
- Mesoscopic level
 - Simulating reactions on spatially extended systems
- **Accurate modelling of catalytic surface reactions**
 - **Complex materials (lattices)**
 - **Complicated reactions**
 - Coverage effects




Graph-Theoretical KMC Approach

- Lattice represented as **graph**
 - Multiple site types
 - Arbitrary connectivity of sites
- State: molecular entity, species, and dentate for every site
 - **Multi-dentate** species allowed
 - **Orientation** explicitly captured





Lattice State Representation




- **Multi-dentate** species modeled (bind to more than one sites)
- Lattice represented as **graph**
- State specifies species & dentation \forall site

Site	Entity	Species	Dentate
1	15	0	1
2	5	0	1
3	16	0	1
⋮	⋮	⋮	⋮
7	21	1	1
18	7	2	1
⋮	⋮	⋮	⋮
24	31	3	1
⋮	⋮	⋮	⋮
31	31	3	2
32	3	0	1





Lattice State Representation




- **Multi-dentate** species modeled (bind to more than one sites)
- Lattice represented as **graph**
- State specifies species & dentation \forall site

Entity	Species	Sites
1	0	4
⋮	⋮	⋮
7	2	18
⋮	⋮	⋮
21	1	7
31	3	24, 31
⋮	⋮	⋮





Elementary Step Representation

- Elementary steps \rightarrow connected graphs
- Subgraph isomorphism** used to
 - identify possible elementary steps
 - map them to lattice processes



Step 178 | $t = 5.83e-013$ sec


Step 179 | $t = 5.83e-013$ sec



Solving the Subgraph Isomorphism


- **Subgraph isomorphism:**

- Create all permutations of N_{sub} out of N_L lattice sites
- Check each permutation



- **Optimizations:**

- Check permutations while constructing (Ullmann 1976)
- Consider only sites within certain distance from entity
 \Rightarrow **localized pattern search**



Lattice Process Lists

- Given lattice state, each **lattice process** fully specified by
 - Type of elementary step
 - Site mapping:

Process	Elem. Step	Sites
1	1	4
2	2	5, 6
3	1	5
4	1	6
5	3	2, 3, 1
⋮	⋮	⋮

Lattice Process ↔ Elementary Step
Which sites are involved in process P_i ?

Entity	#Procs	Processes
1	5	34, 12, 65, 74, 10
2	3	12, 2, 1
3	1	5
4	2	6, 8
5	3	43, 30, 1
⋮	⋮	⋮

Participation Array
Which process entity E_i participates in?

KMC Book-keeping

Process	Elem. Step	Sites
1	1	4
2	2	5, 6
3	1	5
4	1	6
5	3	2, 3, 1
⋮	⋮	⋮

1	1	4
2	2	5, 6
3	1	5
4	1	6
5	3	2, 3, 1
⋮	⋮	⋮

**Lattice
processes**

**Lattice
sites**

**Entities
(adsorbates or
empty sites)**

Entity	#Procs	Processes
1	5	34, 12, 65, 74, 10
2	3	12, 2, 1
3	1	5
4	2	6, 8
5	3	43, 30, 1
⋮	⋮	⋮

1	5	34, 12, 65, 74, 10
2	3	12, 2, 1
3	1	5
4	2	6, 8
5	3	43, 30, 1
⋮	⋮	⋮

Site	Entity	Species	Dentate
1	15	0	1
2	5	0	1
3	16	0	1
⋮	⋮	⋮	⋮
7	21	1	1
18	7	2	1
⋮	⋮	⋮	⋮
24	31	3	1
⋮	⋮	⋮	⋮
31	31	3	2
32	3	0	1

1	15	0	1
2	5	0	1
3	16	0	1
⋮	⋮	⋮	⋮
7	21	1	1
18	7	2	1
⋮	⋮	⋮	⋮
24	31	3	1
⋮	⋮	⋮	⋮
31	31	3	2
32	3	0	1

Entity	Species	Sites
1	0	4
⋮	⋮	⋮
7	2	18
⋮	⋮	⋮
21	1	7
31	3	24, 31
⋮	⋮	⋮


1	0	4
⋮	⋮	⋮
7	2	18
⋮	⋮	⋮
21	1	7
31	3	24, 31
⋮	⋮	⋮

1	0	4
⋮	⋮	⋮
7	2	18
⋮	⋮	⋮
21	1	7
31	3	24, 31
⋮	⋮	⋮

1	0	4
⋮	⋮	⋮
7	2	18
⋮	⋮	⋮
21	1	7
31	3	24, 31
⋮	⋮	⋮


Selecting the Next Event to Happen

- Heap structure:
 - Partially ordered binary tree
 - Each node has **priority** over **all its children nodes**
 - Priority determined by event's execution time
 - **Insert, update, remove** operations reorder the tree so that the **heap property is always satisfied**
 - Next KMC event always found in the top node




Use of Heap Arrays for Queue Construction

- **Heap structures** for the queue of processes to be executed
- Addition, removal, update of a process → **automatic sorting**




Heap Removal Operation

- Assume that process 4 just occurred
- Removal of that process from the heap:




Heap Addition Operation

- Assume we just found another feasible lattice process
- Addition of the 10th process:




Heap Addition Operation

- Assume we just found another feasible lattice process
- Addition of the 10th process:




Heap Addition Operation

- Assume we just found another feasible lattice process
- Addition of the 10th process:




Outline

- Atomistic/Molecular level
 - Calculating rates for elementary events (transition state theory)
- Mesoscopic level
 - Simulating reactions on spatially extended systems
- **Accurate modelling of catalytic surface reactions**
 - Complex materials (lattices)
 - Complicated reactions
 - **Coverage effects**



Non-ideal Adlayers

- Adsorbates on the surface exert attractive/repulsive interactions
 - of the adlayer is not (necessarily) equal to the sum of adsorption energies, *e.g.*:



$$E_{\text{tot}} = 4 \cdot E(\bullet) + 2 \cdot E(\bullet - \bullet) + 2 \cdot E(\bullet \cdots \bullet) + \dots$$

Single body terms
(adsorption energies)

Interaction energies
(2-body and many-body)

- In the general case: cluster expansion

$$\mathcal{H}(\boldsymbol{\sigma}) = \sum_{k=1}^{N_c} \frac{\text{ECI}_k}{\text{GM}_k} \cdot \text{NCE}_k(\boldsymbol{\sigma})$$

Effective cluster interaction

Number of instances of interactions pattern

Graph multiplicity


Patterns represented as graphs, detected using same ideas as for reactions...

Modelling Coverage Effects

- Attractive or repulsive interactions affecting rate, *e.g.*


$$W_{\text{des},i} = A_{\text{des}} \cdot \exp\left(-\frac{E_a - J_{\text{int}} \cdot \sum_{j \in \mathcal{N}_i} \sigma_j}{k_B \cdot T}\right) \cdot \sigma_i$$

Rate increases for
repulsive interactions
($J_{\text{int}} > 0$)



Treating the general case?

Brønsted-Evans-Polanyi Relations



$$E_{\text{fwd}}^\ddagger(\sigma) = E_{\text{fwd},0}^\ddagger + \omega \cdot (\Delta E_{\text{rxn}}(\sigma) - \Delta E_{\text{rxn},0})$$

$$E_{\text{rev}}^\ddagger(\sigma) = E_{\text{rev},0}^\ddagger - (1 - \omega) \cdot (\Delta E_{\text{rxn}}(\sigma) - \Delta E_{\text{rxn},0})$$

$$E_{\text{rev},0}^\ddagger = E_{\text{fwd},0}^\ddagger - \Delta E_{\text{rxn},0}$$

$$\Rightarrow \Delta E_{\text{rxn}}(\sigma) = E_{\text{fwd}}^\ddagger(\sigma) - E_{\text{rev}}^\ddagger(\sigma)$$

... as expected (micro-reversibility)


- Linear correlation between activation and reaction energy
- Captures effect of local reaction environment on rate

Putting It All Together: Algorithm Outline

- Initialize all data structures & populate lattice
- Create a heap of all microscopic processes
- While $t < t_{\text{final}}$
 - Advance time to that of the top process
 - Execute corresponding process:
 - Remove reactants from lattice, and associated processes and energetics
 - Add products into lattice, add new energetic interactions
 - Update event queue:
 - Add to heap all processes in which newly added products can participate
 - Update processes of existing processes if needed (energetic interactions)
- Repeat

Take Home Messages

- Kinetic Monte-Carlo (KMC) simulation:
 - Versatile framework applicable to adsorption/desorption, reaction, diffusion (and other) processes
 - Attracting growing interest in the last few years
- KMC provides unique insight, by bridging
 - molecular scale processes (micro) &
 - observable phenomena (meso, macro)
 - having a dynamic component...



Number of articles published per year containing the keywords "kinetic Monte Carlo" and "catalys" (data from Web of Science)*